ORIGINAL ARTICLE

Diffuse and concentrated recharge evaluation using physical and tracer techniques: results from a semiarid carbonate massif aquifer in southeastern Spain

Francisco J. Alcalá · Yolanda Cantón · Sergio Contreras · Ana Were · Penélope Serrano-Ortiz · Juan Puigdefábregas · Albert Solé-Benet · Emilio Custodio · Francisco Domingo

Received: 26 September 2009/Accepted: 26 March 2010/Published online: 14 April 2010 © Springer-Verlag 2010

Abstract In the high-permeability, semiarid carbonate aquifer in the Sierra de Gádor Mountains (southeastern Spain), some local springs draining shallow perched aquifers were of assistance in assessing applicability of the atmospheric chloride mass balance (CMB) for quantifying total yearly recharge (R_T) by rainfall. Two contrasting hydrological years (October through September) were selected to evaluate the influence of climate on recharge: the average rainfall year 2003–2004, and the unusually dry 2004–2005. Results at small catchment scale were calibrated with estimated daily stand-scale R_T obtained by means of a soil water balance (SWB) of rainfall, using the

F. J. Alcalá Geo-Systems Centre/CVRM, Instituto Superior Técnico, 1049-001 Lisbon, Portugal

F. J. Alcalá (⊠) · J. Puigdefábregas · A. Solé-Benet ·
F. Domingo
Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas, Carretera de Sacramento, s/n, 04120 Almería, Spain
e-mail: fjalcala@eeza.csic.es

Y. Cantón

Departamento de Edafología y Química Agrícola, Universidad de Almería, 04120 Almería, Spain

S. Contreras

Bureau of Economic Geology, Jackson School of Geosciences, University of Texas at Austin, 78758 Austin, TX, USA

A. Were

Department of Hydrology and Geo-environmental Sciences, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands

P. Serrano-Ortiz

Centro Andaluz de Medio Ambiente (CEAMA), 18006 Granada, Spain

actual evapotranspiration measured by the eddy covariance (EC) technique. $R_{\rm T}$ ranged from 0.35 to 0.40 of rainfall in the year, with less than a 5% difference between the CMB and SWB methods in 2003–2004. $R_{\rm T}$ varied from less than 0.05 of rainfall at mid-elevation to 0.20 at high elevation in 2004–2005, with a similar difference between the methods. Diffuse recharge ($R_{\rm D}$) by rainfall was quantified from daily soil water content field data to split $R_{\rm T}$ into $R_{\rm D}$ and the expected concentrated recharge ($R_{\rm C}$) at catchment scale in both hydrological years. $R_{\rm D}$ was 0.16 of rainfall in 2003–2004 and 0.01 in 2004–2005. Under common 1- to 3-day rainfall events, the hydraulic effect of $R_{\rm D}$ is delayed from

P. Serrano-Ortiz
 Department of Ecology, University of Antwerp, 2610 Wilrijk, Belgium

E. Custodio Departament d'Enginyería del Terreny, Cartogràfica i Geofísica, Universitat Politècnica de Catalunya (UPC), 08034 Barcelona, Spain

E. Custodio International Centre for Groundwater Hydrology (FCIHS), Barcelona, Spain

E. Custodio Royal Academy of Sciences of Spain (RAC), 28004 Madrid, Spain

F. Domingo Departamento de Biología Vegetal y Ecología, Universidad de Almería, 04120 Almería, Spain